



LAWRENCE
LIVERMORE
NATIONAL
LABORATORY

Progress report:: NTOF-extracted C rhoR for gas symcap shots

J. Caggiano

November 14, 2013

APS

Denver, CO, United States

November 11, 2013 through November 15, 2013

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.



Measuring Ablator Areal Density using Neutron Time-of-Flight at the National Ignition Facility

**Jac Caggiano, M. Eckart, C. Hagmann, R. Hatarik,
J. McNaney, T. Phillips, D. Casey, D. Sayre, B. Spears, S.
Weber, LLNL**

V. Glebov, J. Knauer, LLE

For the NIF NTOF team

Lawrence Livermore National Laboratory • National Ignition Facility & Photon Science

This work performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344

LLNL-CONF-646165


Outline

- **Motivation**
- **Features in spectrum**
- **Fits to data**
- **Comparison to other carbon rho.R measurements and simulations**
- **Conclude**

The motivation is to measure how much carbon (ablator) is left and to compare to simulations and other measurements




- Initially motivated by seeing what features we could extract from the neutron spectrum as measured by NIF NTOF detectors
- The remaining mass of CH ablator is much larger than in simulations - ~15% range vs. a few percent in simulations
 - Energy not being coupled into ablator as expected
- Measuring this quantity with the NTOF gives a LOS-dependent determination to compare to
 - GRH which is 4π averaged
 - Ross Pair filters which have different directionality





Methodology: Fitting carbon-specific features in the neutron spectrum determine the carbon rho.R

- Two approaches:
 - Fit single scatter cross sections
 - Not preferred because of multiple scattering from higher rho.R shots
 - Fit simulations to data
 - Preferred because simulations include
 - Multiple scattering
- General procedure:
 - Try to fit the data in a region of interest (the $^{12}\text{C}^*[4.44 \text{ MeV}]$ edge), with several simulated spectra assuming golf ball geometry
 - Minimize chi-square parameter in ROI
 - Report CH rho.R
 - Convert to C rho.R by ASSUMING C:H 1:1.38





Methodology: A simple MCNP (neutron transport) model is used to determine spectra to fit to data

- MCNP model:
 - Spherical onion model
 - P0 50 μm (25 for cryo)
 - Dense ablator 50-60 μm
 - DT rho.R 100 mg/cm^2
 - (given amount of gas and shell volume(from sims) this is about right)
 - Inside hohlraum and TMP
 - Temperature = 2.7 keV
 - To match N130505
 - E0 = E0(Tion) (Ballabio)
- Transform through IRF (sensitivity corrected) and fit to the data



How do these features in energy map to features in neutron Time of Flight?


Transformation made for SP LOS
Flight path = 18m




Fitting these features determine
remaining CH rho.R

NTOF diagnostics and NIF Symcap Data


Neutron Time-Of-Flight detectors at NIF measure several implosion variables on many different LOS





Different classes of shots show measureable levels of CH ablator remaining



Simulations overlaid on data show a clear sensitivity to the carbon features




Method 1: Fitting the simulations to a small ROI produce a good measure of the CH ablator areal density







CH rho.R inferred from NTOF SpecSP is 718 ± 71 mg/cm²

Method 2: Using the single scattering cross sections to fit the data produce similar fits and results



Other LOS (alcove) data shows lower CH rho.R than the south pole on all shots



Carbon ρR results are larger on the south pole but average agrees with other diagnostics and simulations




| Carbon ρR | SpecSP $\pm \sim 10\% \text{sys}$ | SpecA | SpecE | LOS avg | GRH | Simulations |
|-----------------|-----------------------------------|----------|----------|------------------|------------|-------------|
| N130503* | <200 | <300 | <300 | | | -- |
| N130505 | 660 \pm 70 | 380(50) | 450(300) | 497(\sim 100) | 520(120) | 485 |
| N130507 | 670 \pm 70 | 380(50) | 450(300) | 500(\sim 100) | 580(120) | 495 |
| N130625 | 500 \pm 60 | 300(50) | 450(300) | 417(\sim 100) | | -- |
| N130628 | 760 \pm 80 | 270(100) | 450(300) | 493(\sim 100) | \sim 600 | -- |

DSR, defined as $N(10-12)/N(13-15)$ is consistent with this picture


| DSR | SpecSP | SpecA | SpecE | Simulations |
|---------|----------|--------|----------|-----------------------|
| N130503 | 0.2(2) | 0.2(1) | 0.45(30) | 0.010 / 0.096 / 0.096 |
| N130505 | 1.5(3) | 1.0(3) | 1.1(4) | 1.04 / 1.01 / 1.00 |
| N130507 | 1.3(3) | 1.1(3) | 1.3(4) | 1.15 / 1.12 / 1.12 |
| N130625 | 1.5(3) | 0.8(3) | 1.1(3) | -- |
| N130628 | 1.45(30) | 0.7(3) | 0.7(3) | 0.69 / 0.68 / 0.69 |

Summary and Future plans

- Carbon rho.R's extracted are in good agreement with GRH
 - Systematic errors of geometry and DT rho.R up to 20%
- Next implement better burn profiles, including a mixed DT/CH region:



D. Casey, S.Weber




B. Spears, S.Weber

NIF





Backup slides

Carbon $^{12}\text{C}(n,n')^{12}\text{C}^*[4.44\text{MeV}]$ feature observed in N121125 and N130510



Compare with N130505 DT symcap where there is very little TT
(CH rho.R = 720 mg/cm²
(10%stat,20%sys)



CH with thin CD layer
T₂, 0.1%D2 100 mg/cm²